



Maternal nutrition during lactation

Authors: Nancy F Butte, PhD, Alison Stuebe, MD, MSc

Section Editors: Steven A Abrams, MD, Kathleen J Motil, MD, PhD

Deputy Editor: Alison G Hoppin, MD

All topics are updated as new evidence becomes available and our [peer review process](#) is complete.

Literature review current through: Jun 2021. | **This topic last updated:** Jun 26, 2020.

INTRODUCTION

A mother's capacity to produce milk of sufficient quantity and quality to support infant growth is resilient and remarkably resistant to nutritional deprivation. However, milk production normally affects maternal body composition and nutritional status, and lactating women have increased nutrient demands.

The changes in maternal nutritional status during lactation, effect of maternal nutrition on milk volume and composition, and nutrient requirements of lactating women are reviewed here. Additional aspects of breastfeeding are discussed separately:

- (See ["Initiation of breastfeeding".](#))
- (See ["Common problems of breastfeeding and weaning".](#))
- (See ["Breastfeeding: Parental education and support".](#))
- (See ["Postpartum contraception: Counseling and methods", section on 'Impact of contraception on breastfeeding'.](#))

EFFECTS OF LACTATION ON THE MOTHER

Lactation is supported partially by mobilization of tissue stores. This, in turn, affects maternal weight and nutritional status.

- **Body weight** – Postpartum weight changes in lactating women are highly variable [1]. Mild, gradual weight loss typically occurs during the first six months postpartum. Excessive weight gain during pregnancy is the most consistent and strongest predictor of

postpartum weight change in most studies [1,2]. Other factors that contribute to weight loss after pregnancy include prepregnancy weight, age, parity, race, smoking, exercise, and return to work outside of the home. (See ["Overview of the postpartum period: Disorders and complications", section on 'Postpartum weight retention'.](#))

Data are conflicting on the effect of breastfeeding on postpartum weight changes. Most studies indicate that lactation has little effect [3-8]. In some reports, weight loss was less in lactating than nonlactating women [2,9-11], whereas other studies report more weight loss in lactating women than nonlactating women [12-15]. A systematic review found little or no association between breastfeeding and weight change, although this seemed to depend on the measurement time points and breastfeeding intensity [16]. However, four out of five studies of high methodologic quality demonstrated a positive association between breastfeeding and weight loss.

- **Lean body mass** – Although some women may lose weight while breastfeeding, the lean body mass of well-nourished women is preserved during the first six months of lactation [17]. This is thought to be due to an adaptive response of protein metabolism when dietary protein is restricted, so that body protein degradation is reduced more than synthesis [18].
- **Vitamins** – Fat-soluble and water-soluble vitamins are secreted into milk. Therefore, the dietary requirement for most vitamins is increased during lactation. A chronically deficient diet will deplete maternal stores of some vitamins. For women consuming a balanced diet, the increased nutrient requirements can be met by the overall increase in food intake. For some restrictive diets, dietary supplements may be necessary. (See ["Vitamins and minerals"](#) below.)
- **Bone mineral loss** – Bone mineral content declines during lactation, and compensatory remineralization occurs after weaning and resumption of menses [19-22]. Changes in calcium, phosphorus, and magnesium homeostasis are independent of maternal dietary intake. Lactation is associated with bone reabsorption, in part due to relatively low estradiol concentrations. A systematic review showed that this bone loss is transient, with recovery depending on the return of menstruation and weaning [23].

Breastfeeding does not increase the risk of bone fractures in the long term. A systematic review that included 11 studies with more than 100,000 women concluded that there is no evidence of an association between breastfeeding and risk of hip, vertebral, or forearm fractures [24].

DETERMINANTS OF MILK VOLUME

Healthy, exclusively breastfeeding women produce approximately 750 to 800 mL per day of milk when lactation is fully established [25]. However, milk volume varies among individuals and can exceed 2000 mL per day in women with hyperlactation or those nursing twins or triplets.

Milk volume is low during the first two days postpartum, increases markedly on days 3 and 4, and then gradually increases to levels seen in full lactation [26]. Milk volume typically decreases later in lactation (eg, beyond six months postpartum), primarily because of weaning. In a cohort of healthy infants, the average (range) of breast milk volumes at 6, 9, and 12 months postpartum were 769 (335 to 1144), 637 (205 to 1185), and 445 (27 to 1154) mL per day, respectively [25]. The wide range of milk volume during late lactation reflects differences in the timing and rate of weaning.

Infant demand — Milk production is determined mostly by infant demand rather than maternal lactation capacity [27]. This is illustrated by the ability of mothers to successfully breastfeed twins or triplets [28]. The infant's demand depends, in turn, on age and rate of weight gain, as well as any medical disorders that affect the infant's metabolic needs or ability to feed efficiently [26,29-31]. Healthy infants who are fed on cue tend to stimulate normal milk volumes and achieve good growth despite individual variation in intervals between feeding [32]. The addition of milk substitutes or solid foods reduces infant demand for breast milk and thus reduces milk production [33,34].

Maternal nutrition — Mild or moderate variations in maternal diet, energy balance, and aerobic exercise generally do not affect milk production [30,35-37].

- **Dieting** – Mild or moderate reductions in maternal energy intake appear to have a limited effect on milk volume [35,38]. In one study, dietary restriction to 1500 kcal per day for one week did not reduce milk production [39]. Similarly, moderate weight loss with or without exercise does not adversely affect lactation [40-42]. In one study, lactating women were randomly assigned to an 11-day program of diet, diet plus exercise, or control at 12 weeks postpartum [41]. Weight loss averaged 1.9, 1.6, and 0.2 kg in the three groups, respectively, while milk volume and composition and infant weight gain were similar during the short study period.

Longer periods of dieting are also well tolerated. In one report, 22 of 33 women who completed a 10-week weight reduction program lost an average of 5 kg [40]. Daily milk production was similar at baseline and at 10 weeks (759 versus 802 mL), and infants gained an average of 21 gm per day. In another study, overweight lactating women were randomly

assigned to reduce energy intake by 500 kcal per day and exercise for 45 minutes four times per week or to maintain their usual diet and limit exercise [43]. Women in the diet and exercise group lost more weight (4.8 versus 0.8 kg) and fat mass (4 versus 0.3 kg) than controls. Infant weight gain was similar, although the statistical power to detect a difference in weight gain was limited [42].

- Severe energy restriction – More extreme nutritional deprivation may have substantial effects on milk volume. In the study cited above, daily milk volume decreased 15 percent when fewer than 1500 kcal per day were consumed [39]. Similarly, some studies of low-income women in resource-limited countries report milk volumes as low as 525 mL/day, while other women produce milk volumes over 800 mL/day, which is similar to average milk volumes in women in resource-rich countries [38,44-50].

Other factors — Other factors that decrease milk volume include incomplete breast emptying and infrequent milk expression [51], as well as maternal smoking, stress, anxiety, fatigue, and illness [52-55]. In addition, use of combined estrogen/progesterone oral contraceptives (COCs) moderately reduces milk volume. In one report that followed lactating women from 6 to 24 weeks postpartum, milk volume decreased by 42 percent in women using COCs, 12 percent in those using progestin-only contraceptives, and 6 percent in women taking no hormonal contraceptives [56]. Nonetheless, infant growth was similar, perhaps because more prolonged or intense feeding periods or supplementary feedings compensated for the reduced milk volume. A systematic review found moderate quality evidence for an effect of COCs on milk volume and overall inconsistent results for other types of hormonal contraceptives [57]. Because there is no firm evidence that hormonal contraceptives interfere with breastfeeding success, an individualized approach seems appropriate [58]. The American College of Obstetrics and Gynecology recommends that "obstetric care providers should discuss any concerns within the context of each woman's desire to breastfeed and her risk of unplanned pregnancy, so that she can make an autonomous and informed decision" [59]. (See "[Postpartum contraception: Counseling and methods](#)", [section on 'Impact of contraception on breastfeeding'](#).)

Skin-to-skin contact between a mother and her preterm infant in the newborn intensive care unit may increase milk production [60,61]. (See "[Breast milk expression for the preterm infant](#)", [section on 'Skin-to-skin contact'](#).)

EFFECTS OF THE MATERNAL DIET ON MILK QUALITY

Lactation requires additional energy and nutrients from the diet. Milk quality is usually sufficient to support infant growth, even when the mother's dietary supply of energy and

nutrients is limited [25]. However, a chronically deficient diet can deplete maternal nutrient stores and adversely affect milk composition.

- **Protein content** – Maternal diet typically does not affect the quantity or quality of milk protein, even in malnourished populations [62-64].
- **Fat content** – The proportion of dietary fat in the maternal diet has minimal effects on the **quantity** of fatty acids in human milk; however, the type of fat consumed by the mother influences the fatty acid composition of the milk [65,66]. As an example, maternal consumption of omega-3 long-chain polyunsaturated fatty acids (n-3 LCPUFA), which include docosahexaenoic acid (DHA), affects concentrations of these fatty acids in breast milk. The relevance of n-3 LCPUFA to infant brain development has been suggested but not confirmed in most clinical studies [67]. (See "[Long-chain polyunsaturated fatty acids \(LCPUFA\) for preterm and term infants](#)".)

Despite lack of definitive evidence, the American Academy of Pediatrics recommends that breastfeeding women take in 200 to 300 mg of n-3 LCPUFA (eg, DHA) to guarantee a sufficient concentration of preformed DHA in the milk [68]. One to two servings of fish per week is adequate to supply this amount of DHA. (See '[Fish intake](#)' below.)

Other variations in the fatty acid content of breast milk may have little clinical relevance. For example, mothers on a low-fat diet produce breast milk with a slightly higher fraction of medium-chain fatty acids compared with mothers consuming a high-fat diet [69]. Consumption of partially hydrogenated fats and oils also influences the fatty acid composition of milk. As an example, linoleic acid concentration was higher in milk from vegan mothers than omnivore controls [70]. However, diet does not appear to affect the cholesterol and phospholipid content of human milk [65].

- **Fat-soluble vitamins** – The concentration of fat-soluble vitamins in milk is reduced in maternal vitamin deficiency and increases following supplementation.

The vitamin D content of breast milk is constitutively low [68,71,72], and it is even lower in mothers with dark skin or other causes of maternal vitamin D deficiency [73]. As a result, all infants require vitamin D supplementation to prevent rickets, and this is particularly important for those whose mothers have vitamin D deficiency. An alternate but rarely used strategy is to administer moderately high doses of vitamin D to the lactating mother (4000 to 6400 international units daily, which is approximately 10 times the recommended intake), which increases the vitamin D content of the breast milk to levels that meet the infant's needs. However, higher doses of vitamin D should be avoided; in one report, ergocalciferol 2500 mcg (100,000 international units daily, which is >150 times the

recommended intake) resulted in toxic levels in milk [74]. (See ["Vitamin D insufficiency and deficiency in children and adolescents", section on 'Maternal vitamin D deficiency'](#) and ["Vitamin D insufficiency and deficiency in children and adolescents", section on 'Prevention in the perinatal period and in infants'.](#))

Vitamin K concentration in breast milk varies with maternal dietary intake but is generally low [75]. As a result, infants are at risk for vitamin K-deficient bleeding, which can be prevented by routine administration of vitamin K at birth. (See ["Overview of vitamin K", section on 'Prevention'.](#))

[Vitamin A](#) content in breast milk also varies with the maternal diet [76,77]. Clinically important vitamin A deficiency is rare in infants in resource-rich countries, in the absence of an underlying malabsorptive disease. In parts of the world in which vitamin A deficiency is endemic, routine vitamin A supplementation is no longer recommended for lactating women, neonates, or infants up to six months of age. This is because studies of vitamin A supplementation have shown no consistent health benefits and may have side effects in young infants. High-dose vitamin A supplementation is recommended in infants and children 6 to 59 months of age in settings where vitamin A deficiency is a public health problem. (See ["Overview of vitamin A", section on 'Universal periodic distribution'.](#))

- **Water-soluble vitamins** – The concentration of water-soluble vitamins in milk also depends upon maternal diet and is reduced with vitamin deficiency. However, the concentration of water-soluble vitamins in milk is regulated so that it does not exceed an upper threshold, even if the mother's intake of these vitamins is very high [78-80]. Vitamin C and [thiamine](#) (vitamin B1) concentrations, for example, remain below approximately 160 mg/L and 200 mcg/L, respectively [78-80].

Vitamin content in milk is reduced in vitamin-deficient states and responds to supplementation [77]. In women with [thiamine](#) deficiency (beriberi), for example, the thiamine content of their milk is low. Vitamin B6 (pyridoxine) concentration in milk also responds rapidly to changes in maternal intake [81].

Folate is preferentially secreted into milk at the expense of maternal reserves. However, milk concentrations are reduced in severe folate deficiency and increase after folate supplementation [82]. [Vitamin B12](#) content in milk is also reduced with deficiency that may occur in vegans, malnourished women, women with latent pernicious anemia [83], and those who have undergone gastric bypass surgery [84]. There are case reports of irreversible developmental delay in exclusively breastfed infants of vitamin B12-deficient

mothers [85]. It may be advisable to monitor maternal vitamin B12 status during lactation, with oral or parenteral supplementation for mothers with vitamin B12 deficiency.

- **Mineral content** – The levels of most inorganic components of human milk are independent of maternal diet or serum concentrations, although intake of some trace minerals affects milk levels. Concentrations of calcium, phosphorus, and magnesium in milk are independent of maternal serum levels and are not substantially influenced by variations in dietary intake [86]. Iron, copper, and zinc levels in human milk are also independent of maternal nutrient status [87-90].

However, breast milk content for some other minerals is associated with maternal nutrient status: In one study, [selenium](#) concentration in milk correlated with maternal plasma concentration, which was influenced by diet [91]. Breast milk iodine concentration also depends upon dietary intake [92]. The primary risk factors for iodine deficiency are a diet without iodized salt or dairy products. Dietary iodine deficiency may also be exacerbated by smoking; iron deficiency; and consumption of large amounts of foods that interfere with the production of thyroid hormones, known as goitrogens, including Brussels sprouts, kale, cabbage, cauliflower, and broccoli [93]. Either iodine deficiency or iodine excess can cause neonatal hypothyroidism [94]. (See "[Clinical features and detection of congenital hypothyroidism](#)", [section on 'Transient congenital hypothyroidism'](#).)

NUTRIENT REQUIREMENTS

Nutritional requirements to support lactation are high, and the requirements for many components are greater in lactating women than in those who are pregnant or in the nonpregnant state ([table 1](#)) [95]. For example, requirements for energy; protein; vitamins A, C, E, B6, and B12; folate; niacin; riboflavin; [thiamine](#); and the minerals iodine, [selenium](#), and zinc are increased in lactating women. In contrast, requirements for vitamins D and K and the minerals calcium, fluoride, magnesium, and phosphorus do not differ between lactating and nonlactating states. The requirement for iron is lower during lactation because of lactational amenorrhea. For women consuming a balanced diet, the increased nutrient requirements can be met by the overall increase in food intake. For some restrictive diets, dietary supplements may be necessary.

Energy — The estimated energy (caloric) requirement during lactation is based on maternal age, weight, height, and physical activity level, as outlined in the table ([table 2](#)) [96], **plus** the following additional energy requirements for milk synthesis:

- From 0 through 6 months postpartum – **330 kcal per day** more than nonlactating women. This is based on the energy cost of exclusive breastfeeding of 500 kcal/day (from an average milk production of 780 mL per day, with an average energy content of 67 kcal/100 mL) [97]. In well-nourished women, the energy cost of lactation is subsidized by mobilization of tissue stores (approximately 170 kcal per day) due to gradual loss of the weight gained during pregnancy during the first six months postpartum.
- From 7 through 12 months of postpartum – **400 kcal per day** more than nonlactating women. The energy cost of lactation is based on an average milk production of 600 mL per day and the same energy content [97]. The calculation for this timeframe does not include a subsidy from mobilization of tissue stores, because it assumes that the mother's weight is stable.

For women with healthy body mass indexes and average heights, this amounts to total energy needs ranging from 2130 to 2730 kcal/day for the first six months of lactation and 2200 to 2800 kcal/day thereafter, depending on maternal age, weight, height, and activity level ([table 2](#)). A woman's energy need also varies with the timing and rate of weaning.

Protein — The recommended dietary allowance (RDA) for protein for the first six months of lactation is 71 grams per day, which is 25 grams more than the requirement for nonlactating women [97]. This RDA is based upon the volume of milk (780 mL per day), the average protein content of milk (1 g/100 mL), and an efficiency of utilization of dietary protein for milk synthesis of 47 percent.

Vitamins and minerals — Recommended intakes for vitamins and minerals during pregnancy and lactation are outlined in the table ([table 1](#)) [86,98,99].

- **Fat-soluble vitamins** – The RDAs for vitamins A and E represent an increase requirement during lactation compared with nonlactating women to compensate for the vitamins secreted into milk. After six months of lactation, the requirement reverts to that of nonlactating women.

The maternal requirements for vitamins D and K are not increased during lactation. However, breast milk does not supply sufficient D and K to meet the infant's needs, and infants require supplementation. Vitamin K supplementation is routinely given to the infant at birth (see "[Overview of vitamin K](#)", [section on 'Prevention'](#)). For the breastfed infant, vitamin D can be provided as supplements or the mother can be supplemented with large doses of vitamin D. (See "[Vitamin D insufficiency and deficiency in children and adolescents](#)", [section on 'Prevention in the perinatal period and in infants'](#).)

- **Water-soluble vitamins** – The RDAs for vitamin C and the B vitamins for lactating women exceed those of nonlactating women ([table 3](#)) [99,100]. These requirements account for the amount of nutrient secreted into milk and allow for metabolic inefficiencies and individual variation.
- **Calcium, phosphorus, and magnesium** – The RDA for calcium during lactation is 1000 mg daily for women 19 years and older and 1300 mg daily for adolescents [86]. These requirements are the same as for nonlactating women, although approximately 200 mg per day of calcium is secreted in milk.

The RDA for calcium is not increased during lactation, because lactation-induced loss in bone mass is not prevented by increased calcium intake and because the loss in bone mass is recovered after weaning [86]. As an example, in one report, loss of bone mass during lactation was similar with calcium supplementation (1 g per day) and placebo [101-103]. Recovery of bone loss occurred with resumption of menses and weaning [104-106]. Similarly, another randomized trial among postpartum women found no difference in bone mineral density with calcium supplementation [107]. Moreover, the calcium content of the breast milk is not affected by maternal calcium intake, because increased bone mobilization and decreased urinary excretion provide the calcium needed for milk production [19,108]. Intestinal calcium absorption does not appear to increase [104,109].

Increased bone resorption and decreased urinary excretion of phosphorus and magnesium are also independent of dietary intake during lactation. Therefore, the dietary recommendations for phosphorus and magnesium are the same as for nonlactating women [110].

- **Trace minerals**

- **Iron** – The RDA for iron during lactation is 9 mg daily for women 19 years and older and 10 mg daily for adolescents [98]. This recommendation is less than the RDA for nonlactating women (18 mg daily) and adolescents (15 mg daily) because of lactation-induced amenorrhea, which reduces iron loss. Thus, postpartum use of a prenatal multivitamin (which provides approximately 30 mg daily of iron) is not necessary for a lactating woman, unless she is known to be iron deficient. A multivitamin that contains less than 30 mg of iron may have fewer gastrointestinal side effects and be better tolerated.
- **Iodine** – In order to support proper infant growth and neurologic development, the RDA for iodine during lactation (290 mcg per day) is higher than pregnancy (220 mcg per day) and prior to pregnancy (150 mcg per day). Several societies recommend that

breastfeeding women take a daily multivitamin or prenatal supplement every day containing 150 mcg of iodine [111].

- **Zinc and selenium** – The RDAs for these minerals are moderately higher during lactation than for nonlactating nonpregnant women to compensate for their secretion into the breast milk [98].

Fish intake — Fish and shellfish contribute high-quality protein and other essential nutrients to the diet. Fish consumption by breastfeeding mothers has been suggested to be beneficial to the infant because fish contain large amounts of omega-3 long-chain polyunsaturated fatty acids (n-3 LCPUFA), including docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA), which are important in brain development. However, available data are insufficient to demonstrate an association between maternal n-3 fatty acid intake during lactation and subsequent cognitive function in offspring. (See "[Long-chain polyunsaturated fatty acids \(LCPUFA\) for preterm and term infants](#)".)

A potential disadvantage of fish intake is exposure to mercury. Nearly all fish and shellfish contain traces of mercury. For most people, the risk from mercury by eating fish and shellfish is not a health concern. Consumption of certain fish and shellfish by breastfeeding women may pose an increased risk to the breastfed infant's developing nervous system since both inorganic and organic mercury are transferred from maternal serum to human milk. The mercury concentrations in human milk are highly variable and tend to decline between colostrum and mature milk [112]. Mercury intake can be minimized by avoiding the intake of predatory fish (eg, pike, marlin, mackerel, tilefish, swordfish).

In considering the potential benefit of n-3 LCPUFA versus the risks of mercury exposure, the American Academy of Pediatrics concluded that the "possible risk from intake of excessive mercury or other contaminants is offset by the neurobehavioral benefits of an adequate DHA intake" [68]. However, some pregnant and postpartum women may have concerns about fish safety, leading them to omit fish consumption altogether to avoid the possibility of mercury exposure [113,114]. Therefore, it is important to provide patients with guidance on choosing safe fish to consume during pregnancy and lactation.

The US Food and Drug Administration and the Environmental Protection Agency issued the following guidelines for fish intake for pregnant women, breastfeeding mothers, and young children [115]. These recommendations allow the mother and her infant to receive some of the potential benefits of eating fish and shellfish, while limiting mercury intake.

- Breastfeeding mothers should eat 8 to 12 ounces a week (two average servings) of fish and shellfish that have lower concentrations of mercury. They include canned light tuna,

salmon, pollock, catfish, and shrimp. Albacore (white) tuna has more mercury than canned light tuna; therefore, consumption of albacore tuna should be limited to 6 ounces (one average meal) per week.

- Breastfeeding mothers should not eat shark, swordfish, king mackerel, or tilefish, because they contain high concentrations of mercury ([table 4](#)).
- Check local advisories about the safety of fish caught by family and friends in local lakes, rivers, and coastal areas. If no advice is available, breastfeeding mothers may eat up to 6 ounces (one average meal) per week of fish caught from local waters but should not consume any other fish during that week.

Potential benefits, risks, and strategies for fish consumption during pregnancy are discussed in detail in a separate topic review. (See "[Fish consumption and marine omega-3 fatty acid supplementation in pregnancy](#)".)

COUNSELING ABOUT COMMON CONCERNS

Special diets — Women with specific dietary needs may have questions about their diets while breastfeeding:

- Dieting for weight loss – After lactation is established, overweight/obese women may restrict their energy intake by 500 kcal/day and undertake aerobic exercise four days/week to promote 0.5 kg/week weight loss without compromising their milk supply [\[116\]](#). Exercise alone, without dieting, does not seem to promote postpartum weight loss. A systematic review concluded that diet or diet plus exercise are effective weight loss strategies for postpartum lactating and nonlactating women [\[117\]](#).

Rarely, a syndrome of "lactation ketoacidosis" has been reported in lactating women, typically associated with restrictive dieting and especially with low carbohydrate intake [\[118,119\]](#). Presenting symptoms and signs include nausea, vomiting, and abdominal pain with a high-anion gap ketoacidosis. It is thought to be a form of fasting ketosis, accelerated by the increased energy needs of a lactating women. (See "[Fasting ketosis and alcoholic ketoacidosis](#)", [section on 'Fasting ketosis'](#).)

- Vegetarian diets – Lactating women on vegetarian diets should be cognizant of the potential risk for mineral, protein, and vitamin deficiencies. Depending on the degree of dietary restriction (eg, ovo- or lactovegetarian), calcium, vitamin D, and [vitamin B12](#) supplements may be required to meet recommended intakes for these nutrients.

- Vegan diet – Women who are healthy but who do **not** eat meat, chicken, fish, or dairy products need to take a vitamin supplement that contains [vitamin B12](#). Most commercially available multivitamins contain an adequate dose of B12.
- Fasting – Short-term fasting does not diminish milk supply but may have a minor effect on milk composition [\[120,121\]](#). Metabolic adaptations during short-term fasting safeguard milk production. Longer-term fasting might adversely affect the nutritional status of lactating women. Therefore, lactating women might be excused from fasting during Ramadan for health reasons [\[121\]](#).
- Food avoidance to prevent atopic disease in offspring – Food avoidance by breastfeeding mothers is not recommended as a means of preventing allergy, even in high-risk children [\[122\]](#). Although both avoidance and ingestion of specific antigens such as peanuts, milk, and eggs during lactation have been hypothesized to reduce the frequency of atopic disease in offspring, the bulk of evidence does not support either approach [\[123\]](#). (See ["Primary prevention of allergic disease: Maternal diet in pregnancy and lactation"](#).)
- Women who have had bariatric surgery – Lactation is usually not adversely affected by bariatric surgery, and breastfeeding should be encouraged. Milk composition of women who have undergone bariatric surgery was shown to be adequate in energy, macronutrients, and [vitamin A](#) [\[124\]](#). Women who breastfeed should continue their vitamin and mineral supplements and should be monitored for nutrient deficiencies, such as [vitamin B12](#) deficiency. (See ["Fertility and pregnancy after bariatric surgery", section on 'Postpartum'](#).)

Alcohol — A small percentage of alcohol is transferred into breast milk. The amount of alcohol considered to be "safe" while breastfeeding is controversial. We suggest that a breastfeeding woman avoid exposing the infant to alcohol by waiting to nurse for two hours after a single serving of alcohol (12 ounces of beer, 5 ounces of wine, or 1.5 ounces of 80-proof liquor). If a woman drinks more than this amount, she should refrain from breastfeeding for an additional two hours for each serving of alcohol [\[125\]](#). It is not necessary to express and discard milk after consuming alcohol, unless the breasts become uncomfortably engorged before enough time has elapsed for the alcohol to leave her system. Heavy alcohol intake can impair judgement and child care abilities and should be avoided, regardless of how the infant is fed. Further details about the pharmacokinetics and effects of alcohol use during lactation are available in the [LactMed](#) database [\[126\]](#).

Caffeine — Most breastfeeding women can drink a moderate amount of caffeine without significant effects on their infants. The American Academy of Pediatrics defines a moderate

intake of caffeine as two to three cups of a caffeinated beverage per day [127]. However, some young infants are sensitive to caffeine and become irritable or have difficulty sleeping, even with small amounts of caffeine. An infant's sensitivity to caffeine usually lessens over time because caffeine clearance is initially slow in newborns but rises to adult clearance levels by three to five months.

Nonnutritive sweeteners — Saccharin, sucralose, and acesulfame potassium have been identified in human breast milk at low levels [126,128,129]. Maternal ingestion of typical quantities probably do not cause adverse effects in the infant, although some experts advise against maternal ingestion of large quantities of these substances. Aspartame is not secreted into human breast milk, even when consumed in quantities far exceeding typical amounts. (See ["Overview of non-nutritive sweeteners", section on 'Specific health outcomes'.](#))

Foods to avoid — Breastfeeding women should avoid fish with high concentrations of mercury ([table 4](#)). (See ["Fish intake"](#) above.)

Otherwise, there are no restrictions of specific foods for the lactating mother. In particular, unpasteurized dairy products or undercooked meats do not have special health risks for lactating women or their infants.

If the mother or infant has a metabolic disease such as phenylketonuria or glucose-6-phosphate dehydrogenase deficiency, special dietary restrictions will apply. (See ["Diagnosis and management of glucose-6-phosphate dehydrogenase \(G6PD\) deficiency", section on 'Management'](#) and ["Overview of phenylketonuria", section on 'Dietary restriction'.](#))

SOCIETY GUIDELINE LINKS

Links to society and government-sponsored guidelines from selected countries and regions around the world are provided separately. (See ["Society guideline links: Breastfeeding and infant nutrition"](#).)

INFORMATION FOR PATIENTS

UpToDate offers two types of patient education materials, "The Basics" and "Beyond the Basics." The Basics patient education pieces are written in plain language, at the 5th to 6th grade reading level, and they answer the four or five key questions a patient might have about a given condition. These articles are best for patients who want a general overview and who prefer short, easy-to-read materials. Beyond the Basics patient education pieces are longer, more

sophisticated, and more detailed. These articles are written at the 10th to 12th grade reading level and are best for patients who want in-depth information and are comfortable with some medical jargon.

Here are the patient education articles that are relevant to this topic. We encourage you to print or e-mail these topics to your patients. (You can also locate patient education articles on a variety of subjects by searching on "patient info" and the keyword(s) of interest.)

- Basics topics (see ["Patient education: Health and nutrition during breastfeeding.\(The Basics\)"](#))
- Beyond the Basics topics (see ["Patient education: Maternal health and nutrition during breastfeeding.\(Beyond the Basics\)"](#))

SUMMARY AND RECOMMENDATIONS

Maternal milk production is dependent upon mobilization of maternal stores and dietary consumption.

- Changes in maternal body composition vary during breastfeeding. Typically, there is a mild gradual weight loss during the first six months postpartum, with preservation of lean body mass. (See ['Effects of lactation on the mother'](#) above.)
- Infant demand is the major factor in determining milk volume. Maternal factors associated with decreased milk volume include maternal stress, anxiety, fatigue, illness, maternal insulin resistance, hormonal oral contraceptives, smoking, and separation from the infant. (See ['Determinants of milk volume'](#) above.)
- Lactation requires additional energy and nutrients in the maternal diet including protein, vitamins A and E, and several other vitamins and minerals ([table 1](#)). (See ['Nutrient requirements'](#) above.)
- The recommended intake for calcium and vitamin D by lactating women are the same as for nonlactating women. This is because maternal bone mineral content, which typically declines during lactation and then rebounds after weaning, is not affected by the mother's intake of these nutrients. Breast milk is low in vitamin D, and supplementation to the infant is required. (See ['Effects of lactation on the mother'](#) above and ['Effects of the maternal diet on milk quality'](#) above and ['Vitamins and minerals'](#) above.)

- Fish consumption by breastfeeding women provides omega-3 long-chain polyunsaturated fatty acids (n-3 LCPUFA), which have been proposed to be beneficial for brain development. However, fish consumption also may pose an increased risk to the breastfed infant's developing nervous system since nearly all fish and shellfish contain traces of mercury and mercury is transferred from maternal serum to human milk. We agree with advisories that suggest women who are pregnant or breastfeeding consume two or three servings per week of fish and shellfish that is high in n-3 LCPUFA and low in mercury, such as canned light tuna, salmon, pollock, tilapia, cod, catfish, and shrimp ([Grade 2C](#)). Shark, swordfish, king mackerel, and tilefish should be avoided because they contain relatively high concentrations of mercury. (See ['Fish intake'](#) above.)
- There are no adverse health effects for the lactating mother or her infant associated with moderate dieting to target gradual weight loss, a balanced vegetarian diet, or short-term fasting. Women consuming a vegan diet or who have undergone bariatric surgery may benefit from nutritional counseling and monitoring to ensure adequate intake of micronutrients, especially [vitamin B12](#). (See ['Special diets'](#) above.)
- Breastfeeding mothers should consume no more than a small amount of alcohol. To minimize transfer of alcohol to the infant, we advise mothers wait at least two hours per serving of alcohol before breastfeeding. Most breastfeeding women can drink a moderate amount of caffeine without significant effects on their infants. (See ['Alcohol'](#) above and ['Caffeine'](#) above.)

Use of UpToDate is subject to the [Subscription and License Agreement](#).

REFERENCES

1. [Butte NF, Hopkinson JM. Body composition changes during lactation are highly variable among women. J Nutr 1998; 128:381S.](#)
2. [Potter S, Hannum S, McFarlin B, et al. Does infant feeding method influence maternal postpartum weight loss? J Am Diet Assoc 1991; 91:441.](#)
3. [Brewer MM, Bates MR, Vannoy LP. Postpartum changes in maternal weight and body fat depots in lactating vs nonlactating women. Am J Clin Nutr 1989; 49:259.](#)
4. [Dugdale AE, Eaton-Evans J. The effect of lactation and other factors on post-partum changes in body-weight and triceps skinfold thickness. Br J Nutr 1989; 61:149.](#)
5. [Ohlin A, Rössner S. Maternal body weight development after pregnancy. Int J Obes 1990; 14:159.](#)

6. [Schauberger CW, Rooney BL, Brimer LM. Factors that influence weight loss in the puerperium. Obstet Gynecol 1992; 79:424.](#)
7. [Parker JD, Abrams B. Differences in postpartum weight retention between black and white mothers. Obstet Gynecol 1993; 81:768.](#)
8. [Greene GW, Smiciklas-Wright H, Scholl TO, Karp RJ. Postpartum weight change: how much of the weight gained in pregnancy will be lost after delivery? Obstet Gynecol 1988; 71:701.](#)
9. [Prasad AS. Zinc deficiency in women, infants and children. J Am Coll Nutr 1996; 15:113.](#)
10. [Neu J. Necrotizing enterocolitis: the search for a unifying pathogenic theory leading to prevention. Pediatr Clin North Am 1996; 43:409.](#)
11. [Rookus MA, Rokebrand P, Burema J, Deurenberg P. The effect of pregnancy on the body mass index 9 months postpartum in 49 women. Int J Obes 1987; 11:609.](#)
12. [DENNIS KJ, BYTHEWAY WR. CHANGES IN BODY WEIGHT AFTER DELIVERY. J Obstet Gynaecol Br Commonw 1965; 72:94.](#)
13. [Kramer FM, Stunkard AJ, Marshall KA, et al. Breast-feeding reduces maternal lower-body fat. J Am Diet Assoc 1993; 93:429.](#)
14. [Dewey KG, Heinig MJ, Nommsen LA. Maternal weight-loss patterns during prolonged lactation. Am J Clin Nutr 1993; 58:162.](#)
15. [Baker JL, Gamborg M, Heitmann BL, et al. Breastfeeding reduces postpartum weight retention. Am J Clin Nutr 2008; 88:1543.](#)
16. [Neville CE, McKinley MC, Holmes VA, et al. The relationship between breastfeeding and postpartum weight change--a systematic review and critical evaluation. Int J Obes \(Lond\) 2014; 38:577.](#)
17. [Motil KJ, Sheng HP, Kertz BL, et al. Lean body mass of well-nourished women is preserved during lactation. Am J Clin Nutr 1998; 67:292.](#)
18. [Motil KJ, Davis TA, Montandon CM, et al. Whole-body protein turnover in the fed state is reduced in response to dietary protein restriction in lactating women. Am J Clin Nutr 1996; 64:32.](#)
19. [Kent GN, Price RI, Gutteridge DH, et al. Human lactation: forearm trabecular bone loss, increased bone turnover, and renal conservation of calcium and inorganic phosphate with recovery of bone mass following weaning. J Bone Miner Res 1990; 5:361.](#)
20. [Lamke B, Brundin J, Moberg P. Changes of bone mineral content during pregnancy and lactation. Acta Obstet Gynecol Scand 1977; 56:217.](#)
21. [Specker BL, Tsang RC, Ho ML. Changes in calcium homeostasis over the first year postpartum: effect of lactation and weaning. Obstet Gynecol 1991; 78:56.](#)

22. [Sowers M, Corton G, Shapiro B, et al. Changes in bone density with lactation. JAMA 1993; 269:3130.](#)
23. [Grizzo FME, Alarcão ACJ, Dell' Agnolo CM, et al. How does women's bone health recover after lactation? A systematic review and meta-analysis. Osteoporos Int 2020; 31:413.](#)
24. Breastfeeding Programs and Policies, Breastfeeding Uptake, and Maternal Health Outcomes in Developed Countries, Feltner C, Weber RP, Stuebe A, Grodensky CA, Orr C, Viswanathan M. (Eds), Agency for Healthcare Research and Quality (US), Rockville (MD) 2018.
25. Institute of Medicine (US) Subcommittee on Lactation. Nutrition during lactation, Committee on Nutritional Status during Pregnancy and Lactation, Food and Nutrition Board, Institute of Medicine, National Academy of Sciences. National Academy Press, Washington, DC, 1991.
26. [Neville MC, Keller R, Seacat J, et al. Studies in human lactation: milk volumes in lactating women during the onset of lactation and full lactation. Am J Clin Nutr 1988; 48:1375.](#)
27. [Macy IG, Hunscher HA, Donelson E, et al. Human milk flow. Am J Dis Child 1930; 39:1186.](#)
28. [Saint L, Maggiore P, Hartmann PE. Yield and nutrient content of milk in eight women breast-feeding twins and one woman breast-feeding triplets. Br J Nutr 1986; 56:49.](#)
29. [Michaelsen KF, Larsen PS, Thomsen BL, Samuelson G. The Copenhagen Cohort Study on Infant Nutrition and Growth: breast-milk intake, human milk macronutrient content, and influencing factors. Am J Clin Nutr 1994; 59:600.](#)
30. [Dewey KG, Heinig MJ, Nommsen LA, Lonnerdal B. Maternal versus infant factors related to breast milk intake and residual milk volume: the DARLING study. Pediatrics 1991; 87:829.](#)
31. [Butte NF, Garza C, Johnson CA, et al. Longitudinal changes in milk composition of mothers delivering preterm and term infants. Early Hum Dev 1984; 9:153.](#)
32. [Butte NF, Garza C, Smith EO, Nichols BL. Human milk intake and growth in exclusively breast-fed infants. J Pediatr 1984; 104:187.](#)
33. [Stuff JE, Nichols BL. Nutrient intake and growth performance of older infants fed human milk. J Pediatr 1989; 115:959.](#)
34. [Dewey KG, Heinig MJ, Nommsen LA, Lönnerdal B. Adequacy of energy intake among breast-fed infants in the DARLING study: relationships to growth velocity, morbidity, and activity levels. Davis Area Research on Lactation, Infant Nutrition and Growth. J Pediatr 1991; 119:538.](#)
35. [Butte NF, Garza C, Stuff JE, et al. Effect of maternal diet and body composition on lactational performance. Am J Clin Nutr 1984; 39:296.](#)

36. [Michaelsen KF. Nutrition and growth during infancy. The Copenhagen Cohort Study. Acta Paediatr Suppl 1997; 420:1.](#)
37. [Dewey KG, Lovelady CA, Nommsen-Rivers LA, et al. A randomized study of the effects of aerobic exercise by lactating women on breast-milk volume and composition. N Engl J Med 1994; 330:449.](#)
38. Prentice A, Paul A, Prentice A, et al. Cross-cultural differences in lactational performance. In: Human Lactation 2: Maternal and Environmental Factors, Hamosh M, Goldman AS (Eds), Plenum Press, New York 1986. p.13.
39. [Strode MA, Dewey KG, Lönnerdal B. Effects of short-term caloric restriction on lactational performance of well-nourished women. Acta Paediatr Scand 1986; 75:222.](#)
40. [Dusdieker LB, Hemingway DL, Stumbo PJ. Is milk production impaired by dieting during lactation? Am J Clin Nutr 1994; 59:833.](#)
41. [McCrary MA, Nommsen-Rivers LA, Molé PA, et al. Randomized trial of the short-term effects of dieting compared with dieting plus aerobic exercise on lactation performance. Am J Clin Nutr 1999; 69:959.](#)
42. [Butte NF. Dieting and exercise in overweight, lactating women. N Engl J Med 2000; 342:502.](#)
43. [Lovelady CA, Garner KE, Moreno KL, Williams JP. The effect of weight loss in overweight, lactating women on the growth of their infants. N Engl J Med 2000; 342:449.](#)
44. [Hennart P, Vis HL. Breast-feeding and post partum amenorrhoea in Central Africa. 1. Milk production in rural areas. J Trop Pediatr 1980; 26:177.](#)
45. [van Steenbergen WM, Kusin JA, Kardjati S, de With C. Energy supplementation in the last trimester of pregnancy in East Java, Indonesia: effect on breast-milk output. Am J Clin Nutr 1989; 50:274.](#)
46. [Brown KH, Akhtar NA, Robertson AD, Ahmed MG. Lactational capacity of marginally nourished mothers: relationships between maternal nutritional status and quantity and proximate composition of milk. Pediatrics 1986; 78:909.](#)
47. [Imong SM, Jackson DA, Wongsawasdi L, et al. Predictors of breast milk intake in rural northern Thailand. J Pediatr Gastroenterol Nutr 1989; 8:359.](#)
48. [de Kanashiro HC, Brown KH, Lopez de Romaña G, et al. Consumption of food and nutrients by infants in Huascar \(Lima\), Peru. Am J Clin Nutr 1990; 52:995.](#)
49. [Orr-Ewing AK, Heywood PF, Coward WA. Longitudinal measurements of breast milk output by a 2H2O tracer technique in rural Papua New Guinean women. Hum Nutr Clin Nutr 1986; 40:451.](#)

50. [Butte NF, Villalpando S, Wong WW, et al. Human milk intake and growth faltering of rural Mesoamerindian infants. Am J Clin Nutr 1992; 55:1109.](#)
51. [Hopkinson JM, Schanler RJ, Garza C. Milk production by mothers of premature infants. Pediatrics 1988; 81:815.](#)
52. [Schanler RJ, Hurst NM. Human milk for the hospitalized preterm infant. Semin Perinatol 1994; 18:476.](#)
53. [Hopkinson JM, Schanler RJ, Fraley JK, Garza C. Milk production by mothers of premature infants: influence of cigarette smoking. Pediatrics 1992; 90:934.](#)
54. [Vio F, Salazar G, Infante C. Smoking during pregnancy and lactation and its effects on breast-milk volume. Am J Clin Nutr 1991; 54:1011.](#)
55. [Andersen AN, Lund-Andersen C, Larsen JF, et al. Suppressed prolactin but normal neurophysin levels in cigarette smoking breast-feeding women. Clin Endocrinol \(Oxf\) 1982; 17:363.](#)
56. [Tankeyoon M, Dusitsin N, Chalapati S, et al. Effects of hormonal contraceptives on milk volume and infant growth. WHO Special Programme of Research, Development and Research Training in Human Reproduction Task force on oral contraceptives. Contraception 1984; 30:505.](#)
57. [Lopez LM, Grey TW, Stuebe AM, et al. Combined hormonal versus nonhormonal versus progestin-only contraception in lactation. Cochrane Database Syst Rev 2015; :CD003988.](#)
58. [Curtis KM, Tepper NK, Jatlaoui TC, et al. U.S. Medical Eligibility Criteria for Contraceptive Use, 2016. MMWR Recomm Rep 2016; 65:1.](#)
59. [ACOG Committee Opinion No. 756: Optimizing Support for Breastfeeding as Part of Obstetric Practice. Obstet Gynecol 2018; 132:e187.](#)
60. [Hurst NM, Valentine CJ, Renfro L, et al. Skin-to-skin holding in the neonatal intensive care unit influences maternal milk volume. J Perinatol 1997; 17:213.](#)
61. [Acuña-Muga J, Ureta-Velasco N, de la Cruz-Bértolo J, et al. Volume of milk obtained in relation to location and circumstances of expression in mothers of very low birth weight infants. J Hum Lact 2014; 30:41.](#)
62. [Lönnerdal B, Forsum E, Gebre-Medhin M, Hambraeus L. Breast milk composition in Ethiopian and Swedish mothers. II. Lactose, nitrogen, and protein contents. Am J Clin Nutr 1976; 29:1134.](#)
63. [Villalpando SF, Butte NF, Wong WW, et al. Lactation performance of rural Mesoamerindians. Eur J Clin Nutr 1992; 46:337.](#)

64. [Sanchez-Pozo A, Lopez Morales J, Izquierdo A, et al. Protein composition of human milk in relation to mothers' weight and socioeconomic status. Hum Nutr Clin Nutr 1987; 41:115.](#)
65. Jensen RG. The Lipids of Human Milk, CRC Press, Boca Raton 1989.
66. [Chappell JE, Francis T, Clandinin MT. Vitamin A and E content of human milk at early stages of lactation. Early Hum Dev 1985; 11:157.](#)
67. [Delgado-Noguera MF, Calvache JA, Bonfill Cosp X, et al. Supplementation with long chain polyunsaturated fatty acids \(LCPUFA\) to breastfeeding mothers for improving child growth and development. Cochrane Database Syst Rev 2015; :CD007901.](#)
68. [Section on Breastfeeding. Breastfeeding and the use of human milk. Pediatrics 2012; 129:e827.](#)
69. [Hachey DL, Silber GH, Wong WW, Garza C. Human lactation. II: Endogenous fatty acid synthesis by the mammary gland. Pediatr Res 1989; 25:63.](#)
70. [Sanders TA, Ellis FR, Dickerson JW. Studies of vegans: the fatty acid composition of plasma choline phosphoglycerides, erythrocytes, adipose tissue, and breast milk, and some indicators of susceptibility to ischemic heart disease in vegans and omnivore controls. Am J Clin Nutr 1978; 31:805.](#)
71. [Misra M, Pacaud D, Petryk A, et al. Vitamin D deficiency in children and its management: review of current knowledge and recommendations. Pediatrics 2008; 122:398.](#)
72. [Henderson A. Vitamin D and the breastfed infant. J Obstet Gynecol Neonatal Nurs 2005; 34:367.](#)
73. [Specker BL, Tsang RC, Hollis BW. Effect of race and diet on human-milk vitamin D and 25-hydroxyvitamin D. Am J Dis Child 1985; 139:1134.](#)
74. [Greer FR, Hollis BW, Napoli JL. High concentrations of vitamin D2 in human milk associated with pharmacologic doses of vitamin D2. J Pediatr 1984; 105:61.](#)
75. [von Kries R, Shearer M, McCarthy PT, et al. Vitamin K1 content of maternal milk: influence of the stage of lactation, lipid composition, and vitamin K1 supplements given to the mother. Pediatr Res 1987; 22:513.](#)
76. [Butte NF, Calloway DH. Evaluation of lactational performance of Navajo women. Am J Clin Nutr 1981; 34:2210.](#)
77. [Daniels L, Gibson RS, Diana A, et al. Micronutrient intakes of lactating mothers and their association with breast milk concentrations and micronutrient adequacy of exclusively breastfed Indonesian infants. Am J Clin Nutr 2019; 110:391.](#)
78. [Bates CJ, Prentice AM, Prentice A, et al. The effect of vitamin C supplementation on lactating women in Keneba, a West African rural community. Int J Vitam Nutr Res 1983;](#)

[53:68.](#)

79. [Byerley LO, Kirksey A. Effects of different levels of vitamin C intake on the vitamin C concentration in human milk and the vitamin C intakes of breast-fed infants. Am J Clin Nutr 1985; 41:665.](#)
80. [PRATT JP, HAMIL BM, MOYER EZ, et al. Metabolism of women during the reproductive cycle. XVIII. The effect of multivitamin supplements on the secretion of B vitamins in human milk. J Nutr 1951; 44:141.](#)
81. [Kang-Yoon SA, Kirksey A, Giacoia G, West K. Vitamin B-6 status of breast-fed neonates: influence of pyridoxine supplementation on mothers and neonates. Am J Clin Nutr 1992; 56:548.](#)
82. [Metz J, Zalusky R, Herbert V. Folic acid binding by serum and milk. Am J Clin Nutr 1968; 21:289.](#)
83. [Johnson PR Jr, Roloff JS. Vitamin B12 deficiency in an infant strictly breast-fed by a mother with latent pernicious anemia. J Pediatr 1982; 100:917.](#)
84. [Celiker MY, Chawla A. Congenital B12 deficiency following maternal gastric bypass. J Perinatol 2009; 29:640.](#)
85. [von Schenck U, Bender-Götze C, Koletzko B. Persistence of neurological damage induced by dietary vitamin B-12 deficiency in infancy. Arch Dis Child 1997; 77:137.](#)
86. Food and Nutrition Board of the Institute of Medicine. Dietary reference intakes for calcium and vitamin D, 2011. Available at: <http://www.nap.edu/catalog/13050/dietary-reference-intakes-for-calcium-and-vitamin-d> (Accessed on June 27, 2016).
87. [Dallman PR. Iron deficiency in the weanling: a nutritional problem on the way to resolution. Acta Paediatr Scand Suppl 1986; 323:59.](#)
88. [Siimes MA, Salmenperä L, Perheentupa J. Exclusive breast-feeding for 9 months: risk of iron deficiency. J Pediatr 1984; 104:196.](#)
89. [Lönnerdal B, Keen CL, Hurley LS. Iron, copper, zinc, and manganese in milk. Annu Rev Nutr 1981; 1:149.](#)
90. [Krebs NF, Reidinger CJ, Hartley S, et al. Zinc supplementation during lactation: effects on maternal status and milk zinc concentrations. Am J Clin Nutr 1995; 61:1030.](#)
91. [Mannan S, Picciano MF. Influence of maternal selenium status on human milk selenium concentration and glutathione peroxidase activity. Am J Clin Nutr 1987; 46:95.](#)
92. [Gushurst CA, Mueller JA, Green JA, Sedor F. Breast milk iodide: reassessment in the 1980s. Pediatrics 1984; 73:354.](#)

93. Centers for Disease Control and Prevention. Breastfeeding: Iodine. Available at: <https://www.cdc.gov/breastfeeding/breastfeeding-special-circumstances/diet-and-micronutrients/iodine.html> (Accessed on March 09, 2020).
94. [Danziger Y, Pertzalan A, Mimouni M. Transient congenital hypothyroidism after topical iodine in pregnancy and lactation. Arch Dis Child 1987; 62:295.](#)
95. [Picciano MF. Nutrient composition of human milk. Pediatr Clin North Am 2001; 48:53.](#)
96. U.S. Department of Agriculture and U.S. Department of Health and Human Services. Dietary Guidelines for Americans, 2020-2025. 9th ed. December 2020. <https://www.dietaryguidelines.gov/> (Accessed on January 13, 2021).
97. DRI, Dietary Reference Intakes: For Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids (Macronutrients). Standing Committee on the Scientific Evaluation of Dietary Reference Intakes, Food, and Nutrition Board, Institute of Medicine. National Academy Press, Washington, DC 2005.
98. DRI, Dietary Reference Intakes: For Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc. Standing Committee on the Scientific Evaluation of Dietary Reference Intakes, Food, and Nutrition Board, Institute of Medicine. National Academy Press, Washington, DC 2000.
99. DRI, Dietary Reference Intakes: For Vitamin C, Vitamin E, Selenium, and Carotenoid. Standing Committee on the Scientific Evaluation of Dietary Reference Intakes, Food, and Nutrition Board, Institute of Medicine. National Academy Press, Washington, DC 2000.
100. Standing Committee on the Scientific Evaluation of Dietary Reference Intakes IoM. Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline. National Academy Press, Washington DC 1998.
101. [Cross NA, Hillman LS, Allen SH, Krause GF. Changes in bone mineral density and markers of bone remodeling during lactation and postweaning in women consuming high amounts of calcium. J Bone Miner Res 1995; 10:1312.](#)
102. [Kalkwarf HJ, Specker BL, Bianchi DC, et al. The effect of calcium supplementation on bone density during lactation and after weaning. N Engl J Med 1997; 337:523.](#)
103. [Prentice A, Jarjou LM, Cole TJ, et al. Calcium requirements of lactating Gambian mothers: effects of a calcium supplement on breast-milk calcium concentration, maternal bone mineral content, and urinary calcium excretion. Am J Clin Nutr 1995; 62:58.](#)
104. [Kalkwarf HJ, Specker BL, Heubi JE, et al. Intestinal calcium absorption of women during lactation and after weaning. Am J Clin Nutr 1996; 63:526.](#)

105. [Sowers M, Eyre D, Hollis BW, et al. Biochemical markers of bone turnover in lactating and nonlactating postpartum women. J Clin Endocrinol Metab 1995; 80:2210.](#)
106. [Hopkinson JM, Butte NF, Ellis K, Smith EO. Lactation delays postpartum bone mineral accretion and temporarily alters its regional distribution in women. J Nutr 2000; 130:777.](#)
107. [Zhang ZQ, Chen YM, Wang RQ, et al. The effects of different levels of calcium supplementation on the bone mineral status of postpartum lactating Chinese women: a 12-month randomised, double-blinded, controlled trial. Br J Nutr 2016; 115:24.](#)
108. [Specker BL, Vieira NE, O'Brien KO, et al. Calcium kinetics in lactating women with low and high calcium intakes. Am J Clin Nutr 1994; 59:593.](#)
109. [Kent GN, Price RI, Gutteridge DH, et al. The efficiency of intestinal calcium absorption is increased in late pregnancy but not in established lactation. Calcif Tissue Int 1991; 48:293.](#)
110. DRI, Dietary reference intakes: For calcium, phosphorus, magnesium, vitamin D, and fluoride. Standing Committee on the Scientific Evaluation of Dietary Reference Intakes, Food, and Nutrition Board, Institute of Medicine. National Academy Press, Washington, DC 1997.
111. [Leung AM, Pearce EN, Braverman LE, Stagnaro-Green A. AAP recommendations on iodine nutrition during pregnancy and lactation. Pediatrics 2014; 134:e1282.](#)
112. [Dórea JG. Exposure to mercury during the first six months via human milk and vaccines: modifying risk factors. Am J Perinatol 2007; 24:387.](#)
113. [Bloomingdale A, Guthrie LB, Price S, et al. A qualitative study of fish consumption during pregnancy. Am J Clin Nutr 2010; 92:1234.](#)
114. [Lando AM, Fein SB, Choinière CJ. Awareness of methylmercury in fish and fish consumption among pregnant and postpartum women and women of childbearing age in the United States. Environ Res 2012; 116:85.](#)
115. Fish: What Pregnant Women and Parents Should Know: Draft Updated Advice by FDA and EPA, June 2014. Available at: <http://www.fda.gov/food/foodborneillnesscontaminants/metals/ucm393070.htm> (Accessed on June 27, 2016).
116. [Lovelady C. Balancing exercise and food intake with lactation to promote post-partum weight loss. Proc Nutr Soc 2011; 70:181.](#)
117. [Amorim Adegboye AR, Linne YM. Diet or exercise, or both, for weight reduction in women after childbirth. Cochrane Database Syst Rev 2013; :CD005627.](#)
118. [Al Alawi AM, Al Flaiti A, Falhammar H. Lactation Ketoacidosis: A Systematic Review of Case Reports. Medicina \(Kaunas\) 2020; 56.](#)
119. [Al Alawi AM, Falhammar H. Lactation ketoacidosis: case presentation and literature review. BMJ Case Rep 2018; 2018.](#)

120. [Neville MC, Sawicki VS, Hay WW Jr. Effects of fasting, elevated plasma glucose and plasma insulin concentrations on milk secretion in women. J Endocrinol 1993; 139:165.](#)
121. [Rakicioğlu N, Samur G, Topçu A, Topçu AA. The effect of Ramadan on maternal nutrition and composition of breast milk. Pediatr Int 2006; 48:278.](#)
122. [Muraro A, Halken S, Arshad SH, et al. EAACI food allergy and anaphylaxis guidelines. Primary prevention of food allergy. Allergy 2014; 69:590.](#)
123. [Pitt TJ, Becker AB, Chan-Yeung M, et al. Reduced risk of peanut sensitization following exposure through breast-feeding and early peanut introduction. J Allergy Clin Immunol 2018; 141:620.](#)
124. [Jans G, Devlieger R, De Preter V, et al. Bariatric Surgery Does Not Appear to Affect Women's Breast-Milk Composition. J Nutr 2018; 148:1096.](#)
125. [Ho E, Collantes A, Kapur BM, et al. Alcohol and breast feeding: calculation of time to zero level in milk. Biol Neonate 2001; 80:219.](#)
126. Drugs and Lactation Database (LactMed). Available at: <https://www.ncbi.nlm.nih.gov/books/NBK501922/> (Accessed on March 09, 2020).
127. [American Academy of Pediatrics Committee on Drugs. Transfer of drugs and other chemicals into human milk. Pediatrics 2001; 108:776.](#)
128. [Rother KI, Sylvetsky AC, Schiffman SS. Non-nutritive sweeteners in breast milk: perspective on potential implications of recent findings. Arch Toxicol 2015; 89:2169.](#)
129. [Sylvetsky AC, Gardner AL, Bauman V, et al. Nonnutritive Sweeteners in Breast Milk. J Toxicol Environ Health A 2015; 78:1029.](#)

Topic 4960 Version 31.0

GRAPHICS

Recommended dietary allowances, or adequate intakes, and tolerable upper limits for adult pregnant and lactating women

	RDAs		ULs for pregnant and lactating women
	Pregnant women*	Lactating women*	
Fat-soluble vitamins			
Vitamin A	770 mcg	1300 mcg	3000 mcg
Vitamin D	600 international units (15 mcg)	600 international units (15 mcg)	4000 international units (100 mcg)
Vitamin E	15 mg	19 mg	1000 mg
Vitamin K [¶]	90 mcg	90 mcg	ND
Water-soluble vitamins			
Vitamin C	85 mg	120 mg	2000 mg
Thiamin	1.4 mg	1.4 mg	ND
Riboflavin	1.4 mg	1.6 mg	ND
Niacin	18 mg	17 mg	35 mg
Vitamin B6	1.9 mg	2 mg	100 mg
Folate	600 mcg	500 mcg	1000 mcg
Vitamin B12	2.6 mcg	2.8 mcg	ND
Minerals			
Calcium	1000 mg	1000 mg	2500 mg
Phosphorus	700 mg	700 mg	4000 mg
Iron	27 mg	9 mg	45 mg
Zinc	11 mg	12 mg	40 mg
Iodine	220 mcg	290 mcg	1100 mcg
Selenium	60 mcg	70 mcg	400 mcg

RDA: recommended dietary allowance; AI: adequate intake; UL: (tolerable) upper limit; ND: not determinable, due to lack of data of adverse effects and concern with regard to lack of ability to handle excess amounts.

* Females over 18 years old.

¶ The requirement for vitamin K is given as an AI rather than an RDA because there was insufficient scientific evidence to calculate the RDA.

Adapted from: *Guidelines for Perinatal Care*, sixth edition (2007); and *Institute of Medicine Dietary Reference Intakes for Calcium and Vitamin D* (2011), which can be accessed via www.nap.edu.

Graphic 60019 Version 9.0

Estimated caloric needs of reproductive-aged women (nonpregnant, nonlactating)

Age (years)	Sedentary* (kcal/day)	Moderately active¶ (kcal/day)	Active ^Δ (kcal/day)
15 to 18	1800	2000	2400
19 to 25	2000	2200	2400
26 to 30	1800	2000	2400
31 to 50	1800	2000	2200
51 to 55	1600	1800	2200

These estimates are based on the estimated energy requirements equations, using reference heights (average) and reference weights (healthy) for each age group. These estimates apply to women who are not pregnant or breastfeeding.

kcal: kilocalories.

* Sedentary means a lifestyle that includes only the physical activity of independent living.

¶ Moderately active means a lifestyle that includes physical activity equivalent to walking approximately 1.5 to 3 miles per day at 3 to 4 miles per hour, in addition to the activities of independent living.

Δ Active means a lifestyle that includes physical activity equivalent to walking more than 3 miles per day at 3 to 4 miles per hour, in addition to the activities of independent living.

Data from: U.S. Department of Health and Human Services and U.S. Department of Agriculture. 2020–2025 Dietary Guidelines for Americans, 9th Edition, December 2020. Available at: <https://www.dietaryguidelines.gov/> (Accessed on January 20, 2021).

Graphic 108793 Version 4.0

Recommended dietary allowances for vitamins during lactation

	RDA for women	Milk secretion	Efficiency	RDA for lactation
Vitamin C	75 mg/d	40 mg/d	85 percent	120 mg/d
Thiamine	1.1 mg/d	0.16 mg/d	0.1 mg/d	1.4 mg/d
Riboflavin	1.1 mg/d	0.3 mg/d	0.1 mg/d	1.6 mg/d
Niacin	14 mg/d	1.4 mg/d	1 mg/d	17 mg/d
Vitamin B6	1.3 mg/d	0.1 mg/d	0.48 mg/d	2 mg/d
Folic acid	400 mcg/d	66 mcg/d	66 mcg/d	500 mcg/d
Vitamin B12	2 mcg/d	0.33 mcg/d		2.8 mcg/d

RDA: recommended dietary allowance.

Data from: *Dietary Reference Intakes. The Essential Guide to Nutrient Requirements*. Otten JJ, Hellwig JP, Meyers LD (Eds), IoM, Washington, DC 2006.

Graphic 62755 Version 6.0

Mercury levels in commercial fish and shellfish

Species	Mercury concentration (PPM; mean)	Number of samples
Fish and shellfish with highest levels of mercury		
King mackerel	0.730	213
Shark	0.988	351
Swordfish	0.976	618
Tilefish (Gulf of Mexico)	1.450	60
Fish and shellfish with lower levels of mercury[¶]		
Anchovies	0.043	40
Butterfish	0.058	89
Catfish	0.049	23
Clam*	ND	6
Cod	0.095	39
Crab ^Δ	0.060	63
Crawfish	0.033	44
Croaker (Atlantic)	0.072	35
Flatfish* [◇]	0.045	23
Haddock (Atlantic)	0.031	4
Hake	0.014	9
Herring	0.044	38
Jacksmelt	0.108	16
Lobster (spiny)	0.09	9
Mackerel (North Atlantic)	0.050	80
Mackerel (Chub, Pacific)	0.088	30
Mullet	0.046	191
Oyster	0.013	38
Perch (ocean)*	ND	6
Pollock	0.041	62
Salmon (canned)*	ND	23
Salmon (fresh/frozen)*	0.014	34
Sardine	0.016	29
Scallop	0.050	66
Shad (American)	0.065	59
Shrimp*	ND	24
Squid	0.070	200
Tilapia*	0.010	9
Trout (freshwater)	0.072	34
Tuna (canned, light)	0.118	347
Whitefish	0.069	28
Whiting	ND	2

Mercury levels of other fish and shellfish ¶		
Bass (saltwater, black, striped) §	0.219	47
Bass (Chilean)	0.386	40
Bluefish	0.337	52
Buffalofish	0.19	4
Carp	0.14	2
Croaker (white, Pacific)	0.287	15
Grouper (all species)	0.465	43
Halibut	0.252	46
Lobster (Northern/American)	0.310	88
Lobster (species unknown)	0.169	16
Mackerel (Spanish, Gulf of Mexico)	0.454	66
Mackerel (Spanish, South Atlantic)	0.182	43
Marlin*	0.485	16
Monkfish	0.180	81
Orange roughy	0.554	49
Perch (freshwater)	0.14	5
Sablefish	0.220	102
Scorpionfish	0.286	78
Sheepshead	0.128	59
Skate	0.137	56
Snapper	0.189	43
Tilefish (Atlantic)	0.144	32
Tuna (canned, Albacore)	0.353	399
Tuna (fresh/frozen, all)	0.383	228
Tuna (fresh/frozen, Albacore)	0.357	26
Tuna (fresh/frozen, Bigeye)	0.639	13
Tuna (fresh/frozen, Skipjack)	0.205	2
Tuna (fresh/frozen, Yellowfin)	0.325	87
Tuna (fresh/frozen, species unknown)	0.414	100
Weakfish (sea trout)	0.256	39

Mercury was measured as total mercury, except for species (*) in which only methylmercury was analyzed.

Source of data: FDA 1990 to 2004, "National Marine Fisheries Service Survey of Trace Elements in the Fishery Resource" Report 1978, "The Occurrence of Mercury in the Fishery Resources of the Gulf of Mexico" Report 2000.

ND: not determinable, since mercury concentration was below the detection level (<0.01 ppm).

¶ The following species have been removed from the tables: Bass (freshwater) - not commercial; Pickerel - not commercial.

Δ Includes blue, king, snow.

◇ Includes flounder, plaice, sole.

§ Includes sea bass/striped bass/rockfish.

Data from: Mercury levels in commercial fish and Shellfish. US Food and Drug Administration. Available at <http://www.fda.gov/Food/FoodSafety/Product-SpecificInformation/Seafood/FoodbornePathogensContaminants/Methylmercury/ucm115644.htm>. Accessed 1/26/2010.

Graphic 63441 Version 6.0

Contributor Disclosures

Nancy F Butte, PhD Nothing to disclose **Alison Stuebe, MD, MSc** Patent Holder: Couplet Care Bassinet [Inventor]. Grant/Research/Clinical Trial Support: Janssen Research and Development [Perinatal depression]. **Steven A Abrams, MD** Grant/Research/Clinical Trial Support: Perrigo Nutrition [Food insecurity; coronavirus disease 2019]. **Kathleen J Motil, MD, PhD** Consultant/Advisory Boards: Acadia. **Alison G Hoppin, MD** Nothing to disclose

Contributor disclosures are reviewed for conflicts of interest by the editorial group. When found, these are addressed by vetting through a multi-level review process, and through requirements for references to be provided to support the content. Appropriately referenced content is required of all authors and must conform to UpToDate standards of evidence.

[Conflict of interest policy](#)

→